

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : C12N 1/21, 15/00, C07K 13/00 C07K 15/28</p>	<p>A1</p>	<p>(11) International Publication Number: WO 94/02590 (43) International Publication Date: 3 February 1994 (03.02.94)</p>
<p>(21) International Application Number: PCT/US93/06733 (22) International Filing Date: 16 July 1993 (16.07.93) (30) Priority data: 07/916,901 20 July 1992 (20.07.92) US (71) Applicant: WAYNE STATE UNIVERSITY [US/US]; 4249 Faculty Administration Bldg., 656 West Kirby, Detroit, MI 48202 (US). (72) Inventors: GRANNEMAN, James, G. ; 5175 Shrewsbury, Troy, MI 48098 (US). LAHNERS, Kristine, N. ; 1713 Hollywood Avenue, Grosse Pointe Woods, MI 48236 (US). RAO, Donald, D. ; 605 West Madison Street #2710, Chicago, IL 60661 (US).</p>		<p>(74) Agent: KOHN, Kenneth, I.; Reising, Ethington, Barnard, Perry & Milton, P.O. Box 4390, Troy, MI 48099 (US). (81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i></p>
<p>(54) Title: β_3-ADRENERGIC RECEPTOR PROTEIN AND DNA ENCODING SAME (57) Abstract Described herein is the β_3-adrenergic receptor protein and DNA which encodes the protein, vectors containing the DNA, host cells transformed with the vectors and methods of using the protein, the DNA and the transformed host cells.</p>		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

-1-

**β_3 -ADRENERGIC RECEPTOR PROTEIN
AND DNA ENCODING SAME**

5

GRANT REFERENCE

The research carried out in connection
with this invention was supported in part by a
10 grant from the NIH (DK37006).

FIELD OF INVENTION

The present invention relates to β_3 -
adrenergic receptor protein, DNA encoding the
15 protein, the genetic elements controlling
expression of the gene, and the use of host cells
transformed with DNA encoding the protein for
screening compounds having utility in modulating
the activity of the β_3 -adrenergic receptor.

20

BACKGROUND OF THE INVENTION

The human β_3 -adrenergic receptor (β_3
receptor) gene was discovered in 1989
(L.J. Emorine et al., Sci. 245, 1989, 1118-1120).
25 The β_3 receptor protein is widely considered to
be a target for agents that will be useful as
human therapeutics (J.R.S. Arch et al., Nature
309, 1984, 163-165), as well as for agents that
beneficially alter the meat and fat content of

SUBSTITUTE SHEET

-2-

feed animals. It has been believed and repeatedly published by those that originally described the β_3 receptor gene that the rodent and human β_3 receptor genes were intronless and
5 that the human gene contained a single exon that encoded a protein of 402 amino acids (Emorine et al., *ibid*; L.J. Emorine et al, *Biochem. Pharmacology* 41, 1991, 853-859; L.J. Emorine et al., *Am. J. Clin. Nutr.* 55 1992, 215S-218S; and
10 C. Nahmias et al. *EMBO Journal* 10, 1991, 3721-3727). DNA constructs have been made that are based upon the assumption that the human β_3 receptor gene contains only 402 amino acids, and these constructs have demonstrated commercial
15 value as reagents for the development of compounds that specifically interact with the β_3 receptor protein.

We have discovered that the assumption that the human β_3 receptor gene contains only one
20 protein-coding block is incorrect. Specifically, we have discovered that the human, rat and mouse β_3 receptor genes contain two protein-coding exons. Thus, the amino acid sequence of the human and mouse β_3 -adrenergic receptor proteins
25 that were previously deduced from genomic DNA are incomplete. Most significantly, we have

SUBSTITUTE SHEET

-3-

discovered that the human β_3 receptor gene is 6 amino acids larger than previously believed. Because we have cloned the human receptor cDNA, we have, for the first time, elucidated the correct amino acid sequence of the human β_3 receptor.

DESCRIPTION OF THE FIGURES

Figure 1. Shows the full coding sequence for the human β_3 -adrenergic receptor sequence and the deduced amino acid.

Figure 2. The structure of the full-length rat β_3 receptor gene.

Figure 3. PCR analysis of rat β_3 receptor cDNA and genomic DNA.

Figure 4. Analysis of rat adipose tissue β_3 receptor mRNA by RNase protection assay. Top: Location of cRNA probe relative to first exon/intron junction. Bottom: Autoradiogram of probe protected by white (WAT), brown (BAT) adipose tissues and liver (LIV). The cRNA probe was fully protected, indicating lack of alternative splicing in these rat tissues.

Figure 5. Comparison of the mouse and human β_3 receptor gene sequences with the homologous sequence of the first exon/intron

SUBSTITUTE SHEET

-4-

junction in the rat gene. Underlined are donor splice signals; the translation termination codons proposed by Emorine et al. (1989, *ibid*) and Nahmias et al. (1991, *ibid*) are in bold.

5 Figure 6. The nucleic acid and deduced amino acid sequences of a partial mouse β_3 receptor cDNA.

 Figure 7. PCR analysis of rat and mouse genomic DNA with cDNA-derived primers.
10 R, rat; M, mouse. See Fig. 3 for location of PCR primers.

 Figure 8. Analysis of β_1 and β_3 receptor mRNA in human omental adipose tissue and in SK-N-MC cells by nuclease protection assay.

15 Figure 9. RNase protection analysis of human β_3 receptor mRNA expressed in SK-N-MC cells.

 Figure 10. Nucleotide and amino acid sequence of a partial human β_3 receptor cDNA (p184). Fig. 10B shows the entire partial
20 sequence and Fig. 10A shows the portion containing the second exon.

 Figure 11. Analysis of β_3 -receptor RNA from human white adipose tissue and SK-N-MC
25 cells.

SUBSTITUTE SHEET

-5-

Figure 12. CHO cells expressing the truncated human β_3 receptor gene make mRNA encoding an unanticipated fusion protein.

Figure 13. Shows a reporter gene
5 construct that expresses rat fat-specific elements.

Figure 14. Shows the construction of the full coding sequence for the human β_3 receptor.

10

SUMMARY OF THE INVENTION

The present invention provides the β_3 -adrenergic receptor protein and DNA which expresses the protein. We have found as
15 described in detail below that previous reports indicating that human β_3 -adrenergic receptor protein is 402 amino acids in length are erroneous, and, in fact, the protein is 408 amino acids in length which provides the basis of the
20 present invention.

The present invention also provides a means for transforming a host cell with a vector containing the DNA which expresses the β_3 -adrenergic receptor and methods of using the
25 transformed host cell for detecting agents, such

SUBSTITUTE SHEET

-6-

as chemical compounds, which affect the activity of the protein.

In another embodiment of the invention, there is provided a means for modifying the DNA which expresses the β_3 -adrenergic receptor protein by site-directed mutagenesis to eliminate a donor splice site to avoid expression of fusion proteins.

Another embodiment of the invention provides oligonucleotide probes which are useful in detecting the presence of mRNA specific for the β_3 -adrenergic receptor protein in cells.

The present invention further provides DNA constructs comprising fat-specific elements of mammalian DNA which expresses β_3 -adrenergic receptor proteins.

Additionally, there is provided novel monoclonal antibodies to the β_3 -adrenergic receptor and fragments thereof.

20

DETAILED DESCRIPTION OF THE INVENTION

In carrying out the work described herein, the following procedures were employed:

General recombinant DNA methods.

Standard cloning techniques used are described by Maniatis et al. (Molecular Cloning: A Laboratory

SUBSTITUTE SHEET

-7-

Manual, 1982, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.). RNA extraction, reverse transcription of tissue RNA and polymerase chain reaction amplification were performed as previously described by Granneman et al. (Endocrinology 130, 1992, 109-114).

Generation of β_3 cDNA probes. Probes for cloning the rat β_3 receptor cDNA and for measurement of tissue mRNA were obtained with the PCR. Brown adipose tissue (BAT) RNA (10 μ g) was reverse-transcribed with a β receptor-specific (Emorine et al., 1989, *ibid*; Kobilka et al., Proc. Natl. Acad. Sci. 84, 1970, 46-50; and Frielle et al., Proc. Natl. Acad. Sci. 84, 1987, 7920-7924) oligonucleotide, primer A, 5'-GCGAATTCGAAGGCACTICIGAAGTCGGGGCTGCGGCAGTA-3', which also contained an EcoRI restriction site on the 5' end. This cDNA was then amplified with primer A and the human β_3 -specific primer 5'-GCGCTGCGCCCGGACAGCTGTGGTCCTGG-3' (Emorine et al., 1989, *ibid*). PCR was performed as described previously by Innis et al. (PCR Protocols, Acad. Press, San Diego, 1990, 54-59). Samples were denatured for 2 min at 94°, annealed, and extended at 72° for 4 mn. Thirty rounds of amplification were performed. One microliter of

SUBSTITUTE SHEET

-8-

this reaction was further amplified, described above, with the β_3 -specific primer described above and a downstream primer, 5'-GCGAATTCGAA-GAAGGGCAGCCAGCAGAG-3', that is common (except for
5 the added EcoRI site) to all β receptors (Emorine et al., 1989, *ibid*; Kobilka et al., 1970, *ibid*; and Frielle et al., 1987, *ibid*). The β_3 receptor PCR product was cloned into the SmaI and EcoRI sites of the plasmid pGEM 3Z (Promega) and
10 sequenced by the dideoxynucleotide chain-termination technique (Sequenase; United States Biochemical Corp). The PCR product was found to be highly homologous to the human β_3 receptor gene (Emorine et al., 1989, *ibid*) and,
15 ultimately, identical to a rat cDNA clone encoding the rat β receptor.

Library construction and screening (rat).

Library construction, screening, and cloning were performed using standard techniques
20 (Maniatis et al., 1982, *ibid*). A cDNA library was constructed in LambdaGEM-4 (Promega) using poly(A)⁺ RNA isolated from BAT of cold-exposed rats. This library contained approximately 3 x 10⁶ recombinants, with an average insert size of
25 1.5 kb. Three hundred thousand recombinants were screened at high stringency (0.03 M NaCl, 3 mM

SUBSTITUTE SHEET

-9-

sodium citrate, pH 7, at 55°) with the cloned rat β_3 PCR product labeled with (32 P)dCTP using random primers (Maniatis et al., *ibid*). Twenty-seven phage were isolated from the amplified library, and two plasmids (p108 and p109) of the same size (about 1.73 kb) were rescued. Sequencing of p108 and p109 from the 5' ends indicated they were identical and truncated with respect to the predicted initiation codon of the human β_3 receptor sequence (Emorine et al., 1989, *ibid*). Screening of the remaining isolates by PCR failed to detect any full-length cDNAs, and primer extension experiments with tissue mRNA suggested that secondary structure, owing to high G-C content, may have limited the ability of the reverse transcriptase to synthesize cDNA through the missing 5' region. Therefore, to obtain the remaining sequence, a Sprague-Dawley rat genomic library (Clontech) was screened with a p108 probe to obtain the rat genomic sequence. The rat β_3 gene was identified by sequencing four hundred forty-four base pairs of genomic sequence that overlapped with p108 β_3 receptor DNA. A full-length clone was then produced by cloning the genomic sequence from bases -104 to +390 (relative to translation initiation) into the

SUBSTITUTE SHEET

-10-

AccI site of p108. Both DNA strands were sequenced by the dideoxy chain-termination technique (Maniatis et al., *ibid*), and no discrepancies were found.

5 Transfection of CHO-k1 cells. The assembled β_3 receptor construct was cloned into pRC/CMV (Invitrogen), an expression vector containing the cytomegalovirus promoter and a neomycin resistance gene. This construct was
10 transfected into CHO-k1 cells using the CaPO_4 method. Stably transfected cells were selected in the presence of Geneticin (800 $\mu\text{g/ml}$) and pooled for further analysis.

Numerous eucaryotic cells can be used.
15 Preferably, these cells will not express any related β adrenergic receptor (i.e., β_1 , β_2 , or β_3 receptors). Examples of such cells include Chinese hamster ovary cells, murine B-82 cells, murine adrenal cortical Y1 cells, xenopus
20 oocytes, or insect Sf cells.

Numerous vectors, some with promoters that are geared to specific cell types can be used. Examples are inducible promoters like mouse mammary tumor virus (MMTV) promoter or
25 metallothionin promoter. Others include retrovirus vectors for gene therapy. Based upon

SUBSTITUTE SHEET

-11-

the information in Example 1 below, numerous variations are possible.

Adenylyl cyclase assay. Adenylyl cyclase activity was determined by the method of Salomon (Adv. Cyclic Nucleotide Res. 10, 1979, 35-55). Culture medium was removed and cells were washed in phosphate-buffered saline and then harvested in 25 mM HEPES (pH 8.0) buffer containing 2 mM $MgCl_2$ and 1 mM EDTA. Cell were homogenized and centrifuged at 48,000 x g for 15 min. to obtain crude membranes. Membrane pellets were resuspended and used directly or frozen at -80° until used. Freezing did not affect activity. Membranes (5-15 μ g of protein) were preincubated at 4°, in a volume of 40 μ l, with the specified drugs for 15 min. Adenylyl cyclase reactions were initiated by addition of substrate mixture and were terminated after 30 min at 30°. BAT membrane adenylyl cyclase activity was determined as previously described (Granneman et al., J. Pharmacol. Exp. Ther. 254, 1990, 508-513, and Granneman et al., J. Pharmacol. Exp. Ther. 256, 1991, 412-425), using membranes from 7-day-old rats. Concentration-response data were analyzed by nonlinear regression analysis with a one-site mass action equation for transfected CHO

SUBSTITUTE SHEET

-12-

cells (Enzfitter, Elsevier Biosoft). A two-site model was used to analyze catecholamine-stimulated adenylyl cyclase in BAT, with the low affinity component representing stimulation by β_3 receptors (Chaudhry et al., Am. Jour. Physiol. 261, 1991, R403-R411).

Tissue mRNA analysis. The size of the β_3 receptor transcripts was determined by Northern blot analysis of rat poly(A)⁺ RNA, as previously described (Maniatis et al., *ibid*; and Granneman et al., Endocrinology 125, 1989, 2328-2335). The cDNA probe used corresponded to bp 228-665 of Fig. 1 and was labeled by random primers. Tissue mRNA distribution experiments were conducted on total RNA with a solution hybridization assay (Maniatis et al., *ibid*; and Granneman et al., Endocrinology 127, 1990, 1596-1601). The radioactive cRNA probe used was transcribed *in vitro* from the cloned β_3 receptor PCR product (p110) with [³²P]CTP, using the T7 promoter. The probe corresponded to bp 746-917 in Fig. 13. Tissue or cellular RNA (6-50 μ g) was co-precipitated with 3×10^4 cpm of the ³²p-labeled cRNA probe. Samples were hybridized at 55° for 12-18 hr and then diluted, and the nonhybridized probe was digested with 300 units

SUBSTITUTE SHEET

-13-

of T-1 ribonuclease for 45 min at 37°. The
[³²P]RNA probe that was protected from RNase
digestion was electrophoretically resolved on a
denaturing polyacrylamide gel containing 8 M
5 urea. The gels were dried and exposed to Kodak
XAR-5 film for 18-72 hr.

Analysis of β_3 receptor mRNA by RNase
protection assay. Rat and human β_3 receptor
mRNAs were analyzed by RNase protection assay
10 using species-specific probes. The rat probe
used (p152) was the BssHII to BglII fragment of
the cloned rat β_3 cDNA cloned into pGEM-7z. This
sequence spans the first exon/intron junction.

Human mRNA was mapped with a β_1
15 receptor probe and two β_3 receptor probes that
were amplified from human genomic DNA. A β_3
receptor (p146) and the β_1 (p145) probes were
amplified by "nested" PCR (Granneman et al.
Molecular Pharmacol. 40, 1991, 895-899) from
20 total nucleic acids using primers based upon the
published sequences (Emorine et al., 1989, *ibid*
and Frielle et al., 1987, *ibid*). The resulting
receptor DNAs were shortened and cloned into
pGEM-7z for the generation of riboprobes. These
25 probes are exact matches of the published
sequences and encode amino acids 178-271 (β_1) and

SUBSTITUTE SHEET

-14-

151 to 223 (β_3). The second human β_3 receptor probe was amplified from genomic DNA (Promega) with a primer set that was designed to amplify a 256 bp DNA fragment which spanned the putative donor splice site. The coding strand primer (HB3G+) was 5'TGCGAATTCTGCCTTCAACCCGCTC 3' and the noncoding strand primer was 5' GCAGGATCCACGGACACATCGCATGCTTCC 3'. Both primers were based upon the published human sequence and contained engineered restriction sites of the 5' ends for cloning into pGEM-7z (p174). The sequence of p174 was an exact match of the published human β_3 receptor gene sequence except for a discrepancy of A for G in the published sequence at bp at 1193 (5, GenBank accession #M29932). This potential discrepancy does not affect the nuclease protection assay because the T-1 ribonuclease used does not cleave at A (J. N. Davidson, The Biochem. of Nucleic Acids, 7th ed., 1972, Academic Press, New York), and no fragments indicative of cleavage at this site were detected.

Cloning of a partial mouse β_3 receptor cDNA.

The mouse β_3 receptor cDNA was obtained from mouse white adipose tissue RNA by reverse

SUBSTITUTE SHEET

-15-

transcription/PCR (Granneman et al., 1992, *ibid*). Reverse transcription of total RNA was performed with the oligonucleotide primer 5' ATTAAAAGGTTTGCATC 3' that was based upon the rat cDNA (Granneman et al., 1991, *ibid*). The resulting cDNA was then amplified by PCR. The coding strand primer was 5' GGACTTTCGCGACGCCT 3' and the noncoding strand primer was 5' GCATCCATGGACGTTGCTTGTC 3', which were also based upon the rat sequence. Samples were denatured at 94°C for 2 min., annealed at 63°C for 1.5 min and extended at 72°C for 2 min for 30 cycles. The resulting PCR product was shortened to 180 bp, cloned into pGEM-7z (p158) and sequenced.

15 PCR analysis of mouse and rat genomic DNA.

To estimate the size of the mouse intron(s), PCR analysis was conducted on mouse and rat genomic DNA. The primer set used was the same that was used above to amplify the mouse cDNA. PCR was carried out for 30 cycles using 1 µg of mouse or rat genomic DNA (Promega) as described above. PCR products were resolved on 1% agarose gel containing ethidium bromide and visualized with ultraviolet light. The identity of these products was verified by Southern blot analysis with an internal probe from the rat cDNA.

SUBSTITUTE SHEET

-16-

Cell culture. SK-N-MC cells were grown in Dulbecco's modified Eagle's medium containing 10% fetal calf serum (Hyclone), penicillin (100,000 units/1) and streptomycin (100 mg/1).

5 Cells were subcultured at a ratio of 1:10 and harvested when about 80% confluent.

Mammalian tissues. Rat tissues were obtained from male Sprague-Dawley rats and mouse tissues were from male outbred mice (Hilltop
10 Labs). Human adipose tissue was obtained with informed consent from surgical specimens.

The discovery of the authentic amino acid sequence of the human β -adrenergic receptor represents a significant improvement in the state
15 of the art with respect to technologies surrounding the β_3 receptor. Specifically, cells expressing the correct amino acid sequence will be most preferable to screen agents for human and animal use. The full length (408 amino acids)
20 human β_3 -adrenergic receptor we have discovered has a pharmacological profile that is different from the truncated (402 amino acids) receptor previously reported. Pindolol derivatives and BRL3744 are partial agonists at both the full
25 length and truncated receptors. However, there is a dramatic difference in the differential

SUBSTITUTE SHEET

-17-

potency of typical β -adrenergic receptor antagonists. Propranolol and alprenolol inhibit the full length receptor with submicromolar potencies, but have been reported to be

5 essentially inactive at the truncated receptor (Emorine et al., 1989, *ibid*). It is difficult to compare the potencies of agonists because of possible differences in receptor reserves, and problems in comparing data from binding and

10 functional measurements. Based on the limited data available, however, it appears that CYP is about 100-fold less potent at the full length receptor than at the truncated receptor.

Pindolol has been reported to have EC_{50} values of

15 150 nM or 1100 nM in truncated β_3 -transfected CHO cells, while we found an EC_{50} of 2800 nM, although a K_i of 84 nM in these cells. Similarly, the EC_{50} for BRL 37344 in truncated β_3 -transfected CHO

cells has been reported to be 6 nM and 180 nM,

20 but we found it to be 840 nM in cells which produce the full-length receptor. It appears that the full-length receptor more closely resembles the "atypical" β -adrenergic receptor found in cardiac, intestinal and adipose tissues

25 and the cloned rat β_3 -receptor than the cloned human truncated β_3 -receptor. Similarities

SUBSTITUTE SHEET

-18-

include 1) a submicromolar affinity for classical
 β -adrenergic receptor antagonists; 2) a
midnanomolar affinity for CYP; 3) activation by
pindolol derivatives with low potency; and 4) a
5 relatively low potency of BRL 37344.

The rodent β_3 receptor is abundantly
expressed only in adipose tissue (Granneman et
al., 1991, *ibid*). In this regard, the original
tissue distribution of the β_3 receptor mRNA
10 described by Emorine et al. (1989, *ibid*) was
erroneous because most of the probe that was used
was derived from the first intron and the exon
sequence used had no homology with the rat
tissues tested. These observations indicate that
15 the β_3 receptor gene contains elements involved
in adipose tissue-specific expression. We have
isolated the rat β_3 receptor gene, and have
identified genetic elements that are likely to be
involved in this phenotypic expression. Tissue-
20 specific enhancers have been identified in the
first intron of several genes (Brooks et al., J.
Biol. Chem. 266, 1991, 7848-7859, and Parmacek et
al., J. Biol. Chem. 265, 1990, 15970-15976). We
have found the sequence within and surrounding
25 one of the inverted repeats in the first intron
of the β_3 receptor gene bears striking homology

SUBSTITUTE SHEET

-19-

with NF-1 (Santaro et al., Natur. 334, 1988, 218-224 and with ARF6 (Graves et al., Mol. Cell. Biol. 12, 1992, 1202-1208). It is anticipated that these sequences are involved in the adipose tissue-specific expression of the β_3 receptor based on recent reports that sequences related to NF-1 and ARF6 are involved in the control of adipose tissue-specific gene expression (Graves et al., 1992, *ibid.* and Genes Dev. 5, 1991, 428-437). The modulation of tissue-specific genes represents a new approach in the treatment of certain diseases and in the generation of agents that produce desirable characteristics in meat-producing animals. For example, agents like Cigilazone that are being developed as antidiabetes therapeutics augment the expression of adipose tissue-specific genes (Kletzien et al., Mol. Pharmacol 41, 1992, 393-398). Efforts to identify novel agents that modify fat-specific gene expression will be facilitated greatly by cell lines expressing readily-detected reporter genes whose transcription is governed by adipose tissue-specific promoter elements. Promoter/reporter gene constructs that are based upon the fat-specific elements within the β_3 receptor gene

SUBSTITUTE SHEET

-20-

represent a novel, useful approach for developing gene-modulating agents.

The structure of DNA sequence of the gene encoding the human β_3 -adrenergic receptor and the amino acid sequence of the receptor was reported by Emorine et al. (1989, *ibid*). The receptor was identified by Emorine et al. as having 402 amino acids, which we have now found to be erroneous, encoded from a single exon. As noted above, we have discovered that the human β_3 -adrenergic gene has two coding exons, and the amino acid sequence for the protein is 408 amino acids in length. We have found that the TAG codon believed to be a termination codon is in a position to contain a human donor splice site (GT) as is more fully detailed below.

The discovery of the donor splice signal in the β_3 -adrenergic receptor gene was initially found in the rat gene. A rat genomic library was screened with the rat β_3 receptor cDNA and isolated a clone containing a 12.1 kb insert. This clone was then subjected to Southern blot analysis using the rat β_3 cDNA as a probe. Digestion of the genomic clone with Xho I revealed prominent bands of 3, 4 and .6 kb that hybridized to the rat β_3 receptor cDNA. Because

SUBSTITUTE SHEET

-21-

the rat β_3 receptor cDNA contains only a single Xho I site, these data suggested the existence of one or more introns in the rat β_3 receptor.

Further analysis utilizing selective cDNA probes
5 suggested the existence of intron(s) near the 3' end of the coding region. The Xho I fragments derived from the genomic clone were then isolated and sequenced.

Shown in figure 2A is a restriction map
10 of the rat β_3 receptor gene and the exon/intron structure of the rat β_3 receptor gene that was deduced by comparison of the genomic sequence with the cDNA (Fig. 2B). "A" shows a map of the rat β_3 receptor gene illustrating the locations
15 of restriction enzyme cleavage sites and the translation initiation (ATG) and termination (TGA) codons. Sequences within this map are contained in the plasmids p111, p108 and p167. "B" shows a schematic representation of the rat
20 β_3 receptor gene, with mature mRNA blocked and the coding sequence filled. E, exon; I, intron. "C" shows a nucleic acid and amino acid sequences of exon/intron junctions of the rat β_3 receptor gene, beginning with Pro³⁷⁴. Underlined are the
25 donor and acceptor splice sites. The inverted repeat that has homology with NF-1 is in bold.

SUBSTITUTE SHEET

-22-

The rat β_3 receptor gene contains three exons and two introns. The first intron interrupts the open reading frame 12 amino acids from the carboxyl end (Fig. 2C). This intron is 394 bp
5 and contains both 5' donor and 3' acceptor splicing signals. The second exon is 68 bp long and encodes the translation termination codon and 28 bp of nontranslated sequence. The second
10 intron is 207 bp long and also contains donor and acceptor splice signals. The final exon contains sequences through the polyadenylation signal as described by Granneman et al. (Molecular Phar.
40, 1991, 895-899).

To further verify that the rat β_3
15 receptor gene contains introns, we performed PCR analysis of rat β_3 receptor cDNA (prepared by reverse transcription of total RNA from adipose tissue) and genomic DNA. PCR primers were
complementary to sequences in the first and third
20 exons (Fig. 3A). The coding strand primer was placed upstream of the first splice junction, whereas the noncoding primer was placed in the third exon. Thus, the expected PCR product
spanned the introns. As expected, this primer
25 set amplified a 845 bp fragment from genomic DNA, as shown in lane 2 of Fig. 3. When tissue cDNA

SUBSTITUTE SHEET

-23-

was used as a template, the product was 246 bp (see lane 1 of Fig 3), as was expected if the primary transcript contained introns which had been removed. No other PCR products were
5 observed, indicating that the β_3 pre-mRNA is not alternatively-placed. To further verify this conclusion, nuclease protection assay was performed on rat β_3 receptor mRNA. The probe used (p152) in this instance was derived from the
10 cloned rat β_3 receptor cDNA and spanned the first exon/intron junction (Fig. 4). If both introns of the rat β_3 receptor are removed by RNA splicing, then tissue mRNA should protect the full (281 nt) complementary probe. However, if
15 the first donor site is not used (i.e., is alternatively spliced), then a fragment of 232 nucleotides would be protected by tissue β_3 receptor mRNA. As shown in Fig. 4, RNA from both white (WAT) and brown (BAT) adipose tissues
20 protected the full probe and no smaller fragments indicative of alternative splicing were observed. As expected, RNA from liver (LIV) failed to protect the β_3 receptor probe indicating that the expression of the gene is adipose tissue-specific
25 (see also Granneman et al., *Endocrinology* 130, 1992, 109-114).

SUBSTITUTE SHEET

-24-

The murine and, as noted above, human β_3 -adrenergic receptor genes have been cloned recently, and both were assumed to lack introns (Emorine et al., 1989, *ibid*, and Nahmias et al., 1991, *ibid*). However, analysis of the genomic sequence alone is not sufficient to decide whether this is so. As shown in Fig. 5, the first exon/intron junction of the rat gene contains the sequence AGGTAG. In the absence of information derived from cDNA, it might be concluded erroneously that the final amino acid is arginine (encoded by AGG) followed by a translation termination codon (TAG). In this regard, we noticed the sequence of the mouse β_3 receptor gene is identical to that of the rat in this region (Nahmias et al., *ibid*). In addition, the human gene also contains the sequence GGTAG in a homologous site, and this sequence has been found to contain a donor splice site (GT), in which case the coding sequence continues, or it could be a termination codon (TAG), as originally deduced.

In order to verify that the mouse gene contains introns, we cloned the relevant region from mouse adipose tissue by reverse transcription of RNA followed by PCR. The

SEQUENCING

-25-

nucleic acid sequence and deduced amino acid sequence of the mouse β_3 receptor cDNA is shown in Fig. 6. The partial cDNA was cloned by reverse transcription of mouse adipose tissue mRNA followed by PCR. Shown is sequence beginning with the codon for Val³⁷⁸. The cDNA exactly matches the genomic sequence reported by Nahmias et al. (1991, *ibid*) until Arg³⁸⁸. The open reading frame continues for 12 more amino acids (Bold, Phe Asp Gly Tyr Glu Gly Ala Arg Pro Phe Pro Thr), which are identical to the rat sequence. The 45 bases of nontranslated sequence in this clone are 71% identical to the non-translated sequence of the rat β_3 receptor cDNA.

To further verify that the mouse β_3 receptor gene contains introns and to estimate their size, PCR analysis of genomic DNA was performed with oligonucleotide primers that were based upon cDNA and were designed to span the intron(s). In the mouse cDNA, there are 208 bp between the primers in this set. Amplification of genomic DNA with this primer set resulted in a PCR product that was about 985 bp, confirming that the mouse gene contains introns and further indicating that the intron(s) present in the

SUBSTITUTE SHEET

-26-

mouse gene are about 120 bp larger than those in the rat gene (Fig. 7).

In order to determine whether the human gene contains introns, we first identified a source of human β_3 receptor mRNA for comparison. In rats, the β_3 adrenergic receptor is expressed abundantly only in adipose tissue, where β_3 receptor mRNA is about 5-7 times more abundant than β_1 receptor mRNA (Granneman et al., Endocrinology 130, 1992, 109-114). We examined mRNA from human subcutaneous and omental adipose tissues by RNase protection assay, and although β_1 receptor mRNA could be readily detected by nuclease protection assay, transcripts encoding the β_3 receptor were absent at the detection limit of the assay (about 4 copies per cell) (Fig. 8). 50 μ g of total RNA was hybridized to human β_1 (p145) and β_3 (p146) receptor probes simultaneously. SK-N-MC cells contain both β_1 and β_3 receptor mRNA, while human omental adipose tissue contains only β_1 receptor transcripts. Right lane shows synthetic human β_3 receptor RNA standards. Thus, although the β_3 receptor does not appear to be expressed in human subcutaneous or omental adipose tissue, we did find that the β_3 receptor is abundantly expressed along with

SUBSTITUTE SHEET

-27-

the β_1 receptor in the human neuroblastoma cell line SK-N-MC. Thus, these cells provide an excellent source for analysis of the human β_3 mRNA (Fig. 8).

5 We mapped the 3' end of the β_3 -adrenergic receptor mRNA from SK-N-MC cells. The probe we used (p174) was derived from human genomic DNA and was designed to span the putative translation termination site/donor splice site
10 (Fig 9, A; see also Fig 5). Referring to Fig. 9, the cRNA probe derived from p146 is complementary to sequence within the first exon of the human β_3 receptor and is fully protected by SK-N-MC mRNA. The cRNA probe derived from p174 is complementary
15 to genomic DNA sequence that spans the putative first exon/intron junction (Fig. 9A). Although SK-N-MC β_3 mRNA protects the full p174 cRNA probe (256 nt), most β_3 transcripts utilize the donor splice signal as indicated by the protected
20 fragment of 194 nt. If the β_3 receptor gene is intronless, then SK-N-MC RNA should fully protect the complementary 256 nucleotide probe. However, if the 5' donor splicing signal contained in the human β_3 receptor pre-mRNA is utilized in the SK-
25 N-MC cells, then cellular RNA should protect exactly 194 nucleotides of the probe. We found

SUBSTITUTE SHEET

-28-

that both 256 and 194 nt of the probe was protected by SK-N-MC RNA (Fig. 9). The ability of SK-N-MC RNA to protect 194 nt of the probe indicated that the splice signals in the human β_3 receptor primary transcript are used by SK-N-MC cells, and thus, the gene contains at least one intron. However, unlike the expression of the rat β_3 receptor gene in adipocytes, the efficiency of splicing was not complete, as indicated by the 256 nt fragment. Thus, about one-fourth of the total β_3 receptor mRNA failed to undergo splicing; and, as originally proposed (Emorine et al., 1989, *ibid*), the translation of the protein would be predicted to terminate at this point. Nevertheless, the great majority of the transcripts were spliced by these cells, and it seemed likely that the human β_3 receptor gene encoded additional amino acids.

To verify whether the spliced human β_3 mRNA encodes additional amino acids, the relevant region of the human β_3 receptor cDNA from SK-N-MC cells was cloned using RACE. Shown in Fig. 10 are the nucleic acid and deduced amino acid sequences of the human β_3 receptor cDNA we obtained. Shown in 10B is the human β_3 receptor cDNA (p184) that was obtained from SK-N-MC cells

SUBSTITUTE SHEET

-29-

using RACE, beginning with the codon for Ala³⁹²
(Fig. 10A). The 5' cDNA sequence of the clone is
identical to the published sequence of the human
gene (Emorine et al., 1991, *ibid*) for 194 bp,
5 then diverges (Bold) exactly at the predicted 5'
donor site. The open reading frame continued for
6 amino acids, followed by 657 bp of
nontranslated sequence. Fig. 10B shows the
complete nucleic acid sequence of p184. Shown in
10 bold is sequence encoding the novel exon(s).
Example 1 sets forth further details of the
cloning of p184.

We also verified that the cDNA sequence
obtained from SK-N-MC cells was in fact expressed
15 in normal human tissues (Fig. 11). Normal human
adipose tissue expresses β_3 receptor mRNA
containing two protein-coding exons. Details of
this experiment are set forth in Example 5
hereof.

20 To Further verify the GT donor splice
site, RNA was obtained from CHO cells that had
been transfected to express the truncated
(encoding 402 amino acids) human β_3 receptor gene
and was subjected to RNase protection analysis
25 with a cRNA probe derived from the human β_3
receptor gene (p174, Fig. 9). CHO cellular RNA

SUBSTITUTE SHEET

-30-

protected both 256 nt and 194 nt of the cRNA probe. The presence of the 194 nt fragment demonstrates that the 5' donor splice signal present in the gene is utilized by CHO cells, and
5 results in the splicing of the first exon with sequences with the expression vector or at the site of DNA integration. Such splicing would be expected to produce a fusion protein, making
10 cells that express such constructs unacceptable for drug screening.

Example 3 set forth below describes means for eliminating fusion proteins by site-directed mutagenesis. The purpose of the site-directed mutagenesis is to alter the codon for
15 gly⁴⁰² so as to eliminate the donor splice signal in order to prevent production of fusion proteins. This modification is important because the splicing of the β_3 receptor premRNA is not complete and can potentially encode both a 402
20 amino acid receptor, as well as a fusion protein.

25

EXAMPLE 1

SUBSTITUTE SHEET

-31-

Cloning of a partial human β_3 receptor
cDNA (p184).

A partial human β_3 receptor cDNA was
cloned by the rapid amplification of cDNA ends
5 (RACE) technique (Frohman et al., 1990, *ibid*).
General cloning techniques used are described in
Maniatis et al., *ibid*. Total RNA (10 μ g) from
SK-N-MC cells was reverse-transcribed as
described previously (Granneman, et al., 1991,
10 *Molecular Pharmacol.* 40, 895-899) with a 17mer
poly T deoxyoligonucleotide primer containing an
engineered XbaI and BamHI restriction sites on the
5' end (5' ACTATAGGGTCTAGAGGATCCGTTTTTTTTTTT-
TTTTTTT 3'). The resulting cDNA was amplified
15 with the human β_3 coding strand was 5'
TGCGAATTCTGCCCTCAACCCGCTC 3. The noncoding
strand primer was 5' ACTATAGGGTCTAGAGGATCCG 3',
which was the adapter sequence of the
primer/adapter oligonucleotide described above.
20 PCR was performed for 30 rounds as follows:
Samples were denatured at 94°C for 2 min.,
annealed at 58°C for 2 min. and extended at 72°C
for 4 min. The resulting products digested with
EcoRI and XbaI, then cloned into pGEM-7z. Twelve
25 recombinants were screened to determine insert
size. Analysis of two clones by RNase protection

SUBSTITUTE SHEET

-32-

assay with the p174 cRNA probe indicated that the 870 bp inserts they contained encoded a human β_3 receptor cDNA. These clones were then analyzed by restriction mapping and dideoxynucleotide sequencing, and were found to be the same. The complete nucleotide sequence of p184 is given in Fig. 10.

EXAMPLE 2

Gene construct Encoding Full-Length
(408 a.a.) Human β_3 -Adrenergic Receptor.

(A) DNA

Such constructs are made as follows: A human β_3 receptor genomic clone is obtained by screening a human genomic library (Clontech) with a radiolabelled probe derived from p184, described above in Example 1. The phage DNA is digested with BglII and BamHI and this 2 kb fragment cloned into pGEM-7z. This construct contains the first exon and part of the first intron of the human β_3 receptor gene. This construct is digested with TaqI (in exon 1) and XbaI (in the vector polylinker). The TaqI to XbaI fragment is removed and the two fragments, containing sequence from XbaI to TaqI (vector/5' gene) and sequence from TaqI to TaqI (in exon 1), is recovered. A three-way ligation is performed

SUBSTITUTE SHEET

-33-

using the fragments just described and the *TaqI* to *XbaI* fragment of p184. The resulting construct contains an open reading frame that encodes the full-length human β_3 receptor. This
5 construct and its preparation are depicted schematically in Fig. 14.

Alternatively, DNA encoding the full-length human β_3 receptor is also obtained by oligonucleotide-directed mutagenesis of truncated
10 (402 amino acid) clones, using commercially-available kits (e.g. Amersham) DNA sequence encoding exon 1 and more than 25 bp of the first intron of the human beta 3 gene (Emorine et al., 1989, *ibid*) is cloned into a M-13 vector, or
15 equivalent, single-stranded vector. Either the coding strand (Fig. 1) or complementary strand may be used. The single-stranded DNA just described is hybridized to an oligonucleotide containing sequence that is complementary to the
20 native genomic strand. The 5' end of the oligonucleotide is complementary to the end of exon 1 to the splice site (Fig. 10A). The next 19 nucleotides begin with G, followed by the codons for ala ser trp gly val ser. The 3' end
25 of the oligonucleotide continues with the sequence at the start of the first intron. For

SUBSTITUTE SHEET

-34-

example: the oligonucleotide 5' CCAGGCTTTGCC-
AACGGCTCGACGGGGCTTCTTGGGGAGTTTCTTAGGTAACGGGGCA-
GAGGGACC 3' (or its complement) is hybridized to
the appropriate single stranded DNA. Useful
5 variants of the oligonucleotide include those
that are somewhat longer or shorter on the 5' or
3' ends. The oligonucleotide is extended with
Klenow polymerase using dCTP α S, and ligated.
Single-stranded DNA is removed, and the native
10 strand is then nicked with NciI and digested with
exonuclease III. The DNA is repolymerized and
ligated, then transformed into host cell (e.g. E.
coli).

(B) Procaryotic and eucaryotic vectors
15 containing DNA described in (A)

The DNA sequence encoding the 408 amino
acid human β_3 receptor protein is first cloned
into an appropriate commercially available vector
for propagation of bacteria. In the present
20 example, the insert described above is cloned
into pGEM-7z. The protein-coding insert is then
shuttled into appropriate mammalian expression
vectors. In this case, we use pRc/CMV
(Invitrogen), an expression vector containing the
25 cytomegalovirus promoter and neomycin resistance
gene. In the case of pRc/CMV, we take advantage

SUBSTITUTE SHEET

-35-

of the HindIII and XbaI sites in the vector to shuttle the insert from pGEM-7z.

(C) Host cells that have been transfected to express the proteins encoded by the DNA constructs described in (B).

(1) Procaryotic cells used to propagate the plasmids are various strains of *E. coli*, including JM109, HB101 and DH5 α . These cells are transformed using standard techniques known to the art.

(2) Eucaryotic cells

Chinese hamster ovary (CHO) cells are transfected with constructs based upon the expression vector pRc/CMV. CHO cells are preferred because they do not natively express any known β receptor subtype. Transfection of CHO cells with DNA constructs is accomplished by the CaPO₄-DNA precipitation method as described in Maniatis et al., *ibid*. To obtain cells that stably express these DNA constructs, transfected cells are selected based upon their resistance to G418, which is conferred by the neomycin-resistance gene contained in pRc/CMV. Cells that survive selective conditions (e.g. 800 μ g/ml G418) are then cloned by limiting dilution. The stable expression of the human β_2 receptor is

SUBSTITUTE SHEET

-36-

verified by (a) the presence of human β_3 ,
receptor mRNA, determined by nuclease protection
assay (see below), and (b) by the stimulation of
adenylyl cyclase (e.g., Granneman et al. *ibid*)
5 with selective β_3 receptor agonists such as BRL
37344 (1 μ M), as well as the stimulation by
isoproterenol (10 μ M) that is resistant to
blockade by CGP 20712A (100 nM).

EXAMPLE 3

10 Construct encoding alternative
(402a.a.) human β_3 receptor wherein Gly⁴⁰² is
degenerate.

DNA constructs encoding the first exon
of the human β_3 receptor in which the codon for
15 glycine⁴⁰² is made degenerate to alter the
sequence GGGTAG so as to eliminate the donor
splice signal is prepared by site-directed
mutagenesis is performed using a commercially
available kit (Amersham) as described in Example
20 2, except that the oligonucleotide is 5'
CCAGGCTTTGCCAACGGCTCGACGG(T/C/A)TAGGTAACCGGGGCAGA
GGGACC 3'. Following this procedure the sequence
GGGTAG will be changed to GGTTAG, GGCTAG, and
GGATAG, respectively.

25

EXAMPLE 4

SUBSTITUTE SHEET

-37-

A method for using the cells described in Example 2C to screen agents that specifically interact (either as agonists or antagonists) with the protein product of such DNA sequences.

5 CHO cells expressing the full length human β_3 receptor are harvested and membranes prepared as described by Granneman et al., *ibid.* Adenylyl cyclase activity is then determined in response to various agents known or thought to
10 interact with the β_3 receptor, using the method of Salomon, *ibid.* Agonists are identified by the ability to increase cyclic AMP generation above basal levels. Antagonists are identified by their ability to decrease adenylyl cyclase
15 activity that is stimulated by 100 nM isoproterenol.

The β_3 receptor is known to increase the formation of cyclic AMP; thus, the interaction of compounds with the recombinant
20 proteins are monitored by changes in cyclic AMP (in whole cells or in cell membranes), or by monitoring the consequences of cyclic AMP formation. There are numerous ways to monitor cyclic AMP, including RIA and fluorescence
25 immunoassay. In addition, the β_3 receptor may activate non-cyclic AMP responses, e.g. calcium

SUBSTITUTE SHEET

-38-

influx. Therefore, there are several functional responses that are the consequence of the activation of these receptors.

Most preferably, these cells will be
5 used to screen compounds that have potential antiobesity, antidiabetes and antispasmodic actions. These cells may also be of benefit in the screening of agents that alter body composition (e.g. repartitioning agents) of meat-
10 producing animals.

EXAMPLE 5

p192 and a method of its use in the detection of human β_3 mRNA.

p184 (see Figs. 10 and 14) contains
15 sequences that are useful in the analysis and detection of mRNA encoding the human β_3 receptor. to obtain one such sequence (p192) the EcoRI to NcoI fragment of p184 was cloned into pGEM-7z. The insert of p192 contains the first 292 bp of
20 p184 and spans exon 1 and exon 2. This construct is used to generate cRNA probes for specific detection of human β_3 receptor mRNA or cDNA, using standard techniques. Shown in Fig. 11 is the use of p192 to detect human β_3 receptor cDNA
25 that had been amplified with PCR. This was performed as follows: RNA from human adipose

SUBSTITUTE SHEET

-39-

tissue and SK-N-MC cells was reverse transcribed with an oligonucleotide primer (5'CAACAGAG-TTGTGCTTCTTGTC 3') that was based upon exon 2 of the cDNA derived from SK-N-MC cells. The
5 resulting cDNA amplified by PCR with this primer and primer HB3G+ (see methods). PCR products were then identified by nuclease protection assay with gene- (p174 - Fig 9) and cDNA-derived (p192) probes (Fig. 11). The fact that human adipose
10 tissue cDNA protects exactly 247 nt of the p192 probe and 194 nt of the p174 probe demonstrate that mRNA corresponding to the novel β_3 receptor cDNA we have cloned from SK-N-MC cells is expressed in normal human adipose tissue (see
15 Fig. 11).

A number of sequences in p184 are useful for diagnosis. These include the PCR primers described above. In general, all sequences that hybridize to either strand of p184
20 are useful. Most preferably, these are sequences (like p192) that can be used to distinguish β_3 receptor mRNA from genomic DNA by DNA amplification techniques (e.g. polymerase chain reaction, for example see Fig. 11) or that can be
25 used to identify or quantify human β_3 receptor mRNA or mRNA splice variants (e.g. ribonuclease

SUBSTITUTE SHEET

-40-

protection assay or RNA hybridization blot analysis (for example Fig. 12). Such sequences are useful in monitoring β_3 receptor gene expression for diagnosis or for development of agents that alter β_3 receptor expression.

EXAMPLE 6

Antibodies that are directed against the amino acid sequence: alanine serine tryptophan glycine valine serine.

10 Polyclonal and monoclonal antibodies are generated against the synthetic peptide by conventional techniques using commercially available services (Chiron, Emeryville, CA). To determine levels of expression of the β_3 receptor, antibodies may be useful in diagnosis to determine levels of expression of the β_3 receptor.

EXAMPLE 7

DNA constructs containing sequences within the introns or 5' flanking regions of the rat β_3 receptor gene (see Fig. 2).

The rat β_3 receptor gene is expressed in a fat-specific fashion (Granneman et al., Molecular Pharmacol., 1991, 40, 895-899). Thus, the gene contains elements that confer fat-specific expression. To obtain the rat β_3

SEQUENCE SHEET

-41-

receptor gene, including the elements that control its expression in fat cells, we screened a rat cDNA library with a rat β_3 cDNA clone. DNA fragments of the gene and cDNA were cloned so as to obtain the DNA sequence of the entire rat gene. The location of the clones obtained are given in Fig. 2 p111 is a 3 kb XhoI to SmaI fragment containing the rat β_3 receptor 5' flanking promoter region); p108 contains the internal 211 bp SmaI to XhoI fragment and p167 contains the 2.6 kb XhoI to SphI fragment, which includes the first and second introns and the second and third exons. Sequence analysis of the first intron of the rat β_3 receptor gene indicates it contains elements involved in fat-specific gene expression.

EXAMPLE 8

Reporter gene constructs that contain elements described in Example 7, that are designed to modify the cellular transcription of the reporter gene.

The following construct (p182) has been made: The NheI to EcoRV fragment of p167, containing the first and second introns of the rat β_3 receptor gene (see 7 above), was cloned into pCAT promoter vector (Promega, Madison WI).

SEQUENCE LISTING

-42-

This construct contains the SV40 promoter, and the introns have been cloned in such a way as to modify the activity of the promoter (Fig. 13). Although we used the chloramphenicol
5 acetyltransferase reporter, there are numerous reporter genes that can be used, e.g. beta galactosidase, luciferase, inter alia). Alternatively, fat-specific elements, especially those in the 5'flanking region contained in p111
10 could be used in the construct.

EXAMPLE 9

Mammalian cells expressing p182.

3T3-F442A cells are stably transfected with constructs described in Example 8 by co-
15 transfecting with pRC/CMV and selection with G418. Stable transformants are identified by increase in reporter gene activity with insulin, and a decrease in reporter gene activity with tumor necrosis factor α .

20 Other appropriate cells into which the construct could be transfected include those that express fat-specific transcription factors or demonstrate the ability to differentiate into an adipocyte phenotype in vitro. Examples of such
25 cells are 3T3-F442A cells, 3T3-L1 cells and RMT

SUBSTITUTE SHEET

-43-

preadipose cells. Included are the nondifferentiated phenotypes of these cells.

Either transient or stable transfections can be used. These can be accomplished by numerous techniques, including CaPO₄, and liposome-mediated transfer (Transfectam, Promega, Madison, Wisconsin) and electroporation. The activity of the reporter gene is monitored by commercially-available kits (e.g. Promega, Madison, Wisconsin).

EXAMPLE 10

A method for using cells described in Example 9 to screen agents for gene-modulating activity.

3T3-F442A cells that have been stably transfected with a β_3 promoter/ β -galactosidase reporter gene construct (p182) are plated in 96-well format. Preadipocytes and differentiated adipocytes are treated with the compound of interest. The activity of the reporter gene will be monitored by the fluorescent product of the Imagene (Molecular probes) β galactosidase substrate with a Cytofluor fluorescence plate reader.

The use of the rat β_3 promoter is not limited to in vitro analysis. The genetic

SUBSTITUTE SHEET

-44-

elements controlling fat-specific expression can be used to target the expression of transgenes to adipocytes of transgenic animals.

EXAMPLE 11

5 Isolation of the introns within the human and murine β_3 receptor gene.

Our discovery of an additional exon in the human and murine β_3 receptor genes allows for the cloning and elucidation of the correct
10 genetic structure of the human and murine β_3 receptor genes. The genetic sequences that intervene the exons in the human gene are obtained by screening a commercially available human genomic library (Clontech) with a
15 radiolabelled BamHI to XbaI fragment of p184 (Fig. 10B). The resulting clone is digested with BamHI, and the bands that hybridize to the full EcoRI to XbaI insert of p184 are gel-isolated and cloned into BamHI linearized, phosphatase-treated
20 pGEM-7z. To obtain the sequences that intervene the novel mouse exons, a commercially-available mouse genomic library (Clontech) is screened with a radiolabelled probe derived from p158, which encodes the novel mouse exons described above.

25 Alternative methods include PCR using oligonucleotides that hybridize to p184 or p158.

SUBSTITUTE SHEET

-45-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Granneman, James G.
Lahners, Kristine N.
Rao, Donald D.

(ii) TITLE OF INVENTION: β 3-ADRENERGIC RECEPTOR PROTEIN AND DNA
ENCODING SAME

(iii) NUMBER OF SEQUENCES: 9

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: REISING, ETHINGTON, BARNARD, PERRY & MILTON
(B) STREET: 201 W. Big Beaver - Ste. 400; P.O. Box 4390
(C) CITY: Troy
(D) STATE: Michigan
(E) COUNTRY: USA
(F) ZIP: 48099

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: US 916,901
(B) FILING DATE: 20-JUL-1992
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Kohn, Kenneth I.
(B) REGISTRATION NUMBER: 30,955
(C) REFERENCE/DOCKET NUMBER: P-324(WSU)

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (313) 689-3554

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1227 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(ix) FEATURE:

(A) NAME/KEY: CDS
(B) LOCATION: 1..1224

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

ATG GCT CCG TGG CCT CAC GAG AAC AGC TCT CTT GCC CCA TGG CCG GAC 48
Met Ala Pro Trp Pro His Glu Asn Ser Ser Leu Ala Pro Trp Pro Asp
1 5 10 15

SUBSTITUTE SHEET

-46-

CTC	CCC	ACC	CTG	GCG	CCC	AAT	ACC	GCC	AAC	ACC	AGT	GGG	CTG	CCA	GGG	96
Leu	Pro	Thr	Leu	Ala	Pro	Asn	Thr	Ala	Asn	Thr	Ser	Gly	Leu	Pro	Gly	
			20					25					30			
GTT	CCG	TGG	GAG	GCG	GCC	CTA	GCC	GGG	GCC	CTG	CTG	GCG	CTG	GCG	GTG	144
Val	Pro	Trp	Glu	Ala	Ala	Leu	Ala	Gly	Ala	Leu	Leu	Ala	Leu	Ala	Val	
		35					40					45				
CTG	GCC	ACC	GTG	GGA	GGC	AAC	CTG	CTG	GTC	ATC	GTG	GCC	ATC	GCC	TGG	192
Leu	Ala	Thr	Val	Gly	Gly	Asn	Leu	Leu	Val	Ile	Val	Ala	Ile	Ala	Trp	
		50				55					60					
ACT	CCG	AGA	CTC	CAG	ACC	ATG	ACC	AAC	GTG	TTC	GTG	ACT	TCG	CTG	GCC	240
Thr	Pro	Arg	Leu	Gln	Thr	Met	Thr	Asn	Val	Phe	Val	Thr	Ser	Leu	Ala	
		65			70					75					80	
GCA	GCC	GAC	CTG	GTG	ATG	GGA	CTC	CTG	GTG	GTG	CCG	CCG	GCG	GCC	ACC	288
Ala	Ala	Asp	Leu	Val	Met	Gly	Leu	Leu	Val	Val	Pro	Pro	Ala	Ala	Thr	
				85					90					95		
TTG	GCG	CTG	ACT	GGC	CAC	TGG	CCG	TTG	GGC	GCC	ACT	GGC	TGC	GAG	CTG	336
Leu	Ala	Leu	Thr	Gly	His	Trp	Pro	Leu	Gly	Ala	Thr	Gly	Cys	Glu	Leu	
			100					105					110			
TGG	ACC	TCG	GTG	GAC	GTG	CTG	TGT	GTG	ACC	GCC	AGC	ATC	GAA	ACC	CTG	384
Trp	Thr	Ser	Val	Asp	Val	Leu	Cys	Val	Thr	Ala	Ser	Ile	Glu	Thr	Leu	
		115					120					125				
TGC	GCC	CTG	GCC	GTG	GAC	CGC	TAC	CTG	GCT	GTG	ACC	AAC	CCG	CTG	CGT	432
Cys	Ala	Leu	Ala	Val	Asp	Arg	Tyr	Leu	Ala	Val	Thr	Asn	Pro	Leu	Arg	
		130				135						140				
TAC	GGC	GCA	CTG	GTC	ACC	AAG	CGC	TGC	GCC	CGG	ACA	GCT	GTG	GTC	CTG	480
Tyr	Gly	Ala	Leu	Val	Thr	Lys	Arg	Cys	Ala	Arg	Thr	Ala	Val	Val	Leu	
		145			150					155					160	
GTG	TGG	GTC	GTG	TCG	GCC	GCG	GTG	TCG	TTT	GCG	CCC	ATC	ATG	AGC	CAG	528
Val	Trp	Val	Val	Ser	Ala	Ala	Val	Ser	Phe	Ala	Pro	Ile	Met	Ser	Gln	
				165					170					175		
TGG	TGG	CGC	GTA	GGG	GCC	GAC	GCC	GAG	GCG	CAG	CGC	TGC	CAC	TCC	AAC	576
Trp	Trp	Arg	Val	Gly	Ala	Asp	Ala	Glu	Ala	Gln	Arg	Cys	His	Ser	Asn	
			180					185					190			
CCG	CGC	TGC	TGT	GCC	TTC	GCC	TCC	AAC	ATG	CCC	TAC	GTG	CTG	CTG	TCC	624
Pro	Arg	Cys	Cys	Ala	Phe	Ala	Ser	Asn	Met	Pro	Tyr	Val	Leu	Leu	Ser	
		195					200					205				
TCC	TCC	GTC	TCC	TTC	TAC	CTT	CCT	CTT	CTC	GTG	ATG	CTC	TTC	GTC	TAC	672
Ser	Ser	Val	Ser	Phe	Tyr	Leu	Pro	Leu	Leu	Val	Met	Leu	Phe	Val	Tyr	
		210				215					220					
GCG	CGG	GTT	TTC	GTG	GTG	GCT	ACG	CGC	CAG	CTG	CGC	TTG	CTG	CGC	GGG	720
Ala	Arg	Val	Phe	Val	Val	Ala	Thr	Arg	Gln	Leu	Arg	Leu	Leu	Arg	Gly	
		225			230					235					240	
GAG	CTG	GGC	CGC	TTT	CCG	CCC	GAG	GAG	TCT	CCG	CCG	GCG	CCG	TCG	CGC	768
Glu	Leu	Gly	Arg	Phe	Pro	Pro	Glu	Glu	Ser	Pro	Pro	Ala	Pro	Ser	Arg	
				245					250					255		
TCT	CTG	GCC	CCG	GCC	CCG	GTG	GGG	ACG	TGC	GCT	CCG	CCC	GAA	GGG	GTG	816
Ser	Leu	Ala	Pro	Ala	Pro	Val	Gly	Thr	Cys	Ala	Pro	Pro	Glu	Gly	Val	
			260					265					270			

SUBSTITUTE SHEET

-47-

CCC GCC TGC GGC CGG CGG CCC GCG CGC CTC CTG CCT CTC CGG GAA CAC	864
Pro Ala Cys Gly Arg Arg Pro Ala Arg Leu Leu Pro Leu Arg Glu His	
275 280 285	
CGG GCC CTG TGC ACC TTG GGT CTC ATC ATG GGC ACC TTC ACT CTC TGC	912
Arg Ala Leu Cys Thr Leu Gly Leu Ile Met Gly Thr Phe Thr Leu Cys	
290 295 300	
TGG TTG CCC TTC TTT CTG GCC AAC GTG CTG CGC GCC CTG GGG GGC CCC	960
Trp Leu Pro Phe Phe Leu Ala Asn Val Leu Arg Ala Leu Gly Gly Pro	
305 310 315 320	
TCT CTA GTC CCG GGC CCG GCT TTC CTT GCC CTG AAC TGG CTA GGT TAT	1008
Ser Leu Val Pro Gly Pro Ala Phe Leu Ala Leu Asn Trp Leu Gly Tyr	
325 330 335	
GCC AAT TCT GCC TTC AAC CCG CTC ATC TAC TGC CGC AGC CCG GAC TTT	1056
Ala Asn Ser Ala Phe Asn Pro Leu Ile Tyr Cys Arg Ser Pro Asp Phe	
340 345 350	
CGC AGC GCC TTC CGC CGT CTT CTG TGC CGC TGC GGC CGT CGC CTG CCT	1104
Arg Ser Ala Phe Arg Arg Leu Leu Cys Arg Cys Gly Arg Arg Leu Pro	
355 360 365	
CCG GAG CCC TGC GCC GCC GCC CGC CCG GCC CTC TTC CCC TCG GGC GTT	1152
Pro Glu Pro Cys Ala Ala Ala Arg Pro Ala Leu Phe Pro Ser Gly Val	
370 375 380	
CCT GCG GCC CGG AGC AGC CCA GCG CAG CCC AGG CTT TGC CAA CGG CTC	1200
Pro Ala Ala Arg Ser Ser Pro Ala Gln Pro Arg Leu Cys Gln Arg Leu	
385 390 395 400	
GAC GGG GCT TCT TGG GGA GTT TCT TAG	1227
Asp Gly Ala Ser Trp Gly Val Ser	
405	

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 408 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Ala	Pro	Trp	Pro	His	Glu	Asn	Ser	Ser	Leu	Ala	Pro	Trp	Pro	Asp
1				5					10					15	
Leu	Pro	Thr	Leu	Ala	Pro	Asn	Thr	Ala	Asn	Thr	Ser	Gly	Leu	Pro	Gly
			20					25					30		
Val	Pro	Trp	Glu	Ala	Ala	Leu	Ala	Gly	Ala	Leu	Leu	Ala	Leu	Ala	Val
			35				40					45			
Leu	Ala	Thr	Val	Gly	Gly	Asn	Leu	Leu	Val	Ile	Val	Ala	Ile	Ala	Trp
	50					55				60					
Thr	Pro	Arg	Leu	Gln	Thr	Met	Thr	Asn	Val	Phe	Val	Thr	Ser	Leu	Ala
65					70					75				80	

SUBSTITUTE SHEET

-48-

Ala Ala Asp Leu Val Met Gly Leu Leu Val Val Pro Pro Ala Ala Thr
 85 90 95
 Leu Ala Leu Thr Gly His Trp Pro Leu Gly Ala Thr Gly Cys Glu Leu
 100 105 110
 Trp Thr Ser Val Asp Val Leu Cys Val Thr Ala Ser Ile Glu Thr Leu
 115 120 125
 Cys Ala Leu Ala Val Asp Arg Tyr Leu Ala Val Thr Asn Pro Leu Arg
 130 135 140
 Tyr Gly Ala Leu Val Thr Lys Arg Cys Ala Arg Thr Ala Val Val Leu
 145 150 155 160
 Val Trp Val Val Ser Ala Ala Val Ser Phe Ala Pro Ile Met Ser Gln
 165 170 175
 Trp Trp Arg Val Gly Ala Asp Ala Glu Ala Gln Arg Cys His Ser Asn
 180 185 190
 Pro Arg Cys Cys Ala Phe Ala Ser Asn Met Pro Tyr Val Leu Leu Ser
 195 200 205
 Ser Ser Val Ser Phe Tyr Leu Pro Leu Leu Val Met Leu Phe Val Tyr
 210 215 220
 Ala Arg Val Phe Val Val Ala Thr Arg Gln Leu Arg Leu Leu Arg Gly
 225 230 235 240
 Glu Leu Gly Arg Phe Pro Pro Glu Glu Ser Pro Pro Ala Pro Ser Arg
 245 250 255
 Ser Leu Ala Pro Ala Pro Val Gly Thr Cys Ala Pro Pro Glu Gly Val
 260 265 270
 Pro Ala Cys Gly Arg Arg Pro Ala Arg Leu Leu Pro Leu Arg Glu His
 275 280 285
 Arg Ala Leu Cys Thr Leu Gly Leu Ile Met Gly Thr Phe Thr Leu Cys
 290 295 300
 Trp Leu Pro Phe Phe Leu Ala Asn Val Leu Arg Ala Leu Gly Gly Pro
 305 310 315 320
 Ser Leu Val Pro Gly Pro Ala Phe Leu Ala Leu Asn Trp Leu Gly Tyr
 325 330 335
 Ala Asn Ser Ala Phe Asn Pro Leu Ile Tyr Cys Arg Ser Pro Asp Phe
 340 345 350
 Arg Ser Ala Phe Arg Arg Leu Leu Cys Arg Cys Gly Arg Arg Leu Pro
 355 360 365
 Pro Glu Pro Cys Ala Ala Ala Arg Pro Ala Leu Phe Pro Ser Gly Val
 370 375 380
 Pro Ala Ala Arg Ser Ser Pro Ala Gln Pro Arg Leu Cys Gln Arg Leu
 385 390 395 400
 Asp Gly Ala Ser Trp Gly Val Ser
 405

SEQUENCE SHEET

-49-

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 870 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 1..195

(ix) FEATURE:

- (A) NAME/KEY: exon
- (B) LOCATION: 196..870

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..215

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

AAT TCT GCC TTC AAC CCG CTC ATC TAC TGC CGC AGC CCG GAC TTT CGC	48
Asn Ser Ala Phe Asn Pro Leu Ile Tyr Cys Arg Ser Pro Asp Phe Arg	
1 5 10 15	
AGC GCC TTC CGC CGT CTT CTG TGC CGC TGC GGC CGT CGC CTG CCT CCG	96
Ser Ala Phe Arg Arg Leu Leu Cys Arg Cys Gly Arg Arg Leu Pro Pro	
20 25 30	
GAG CCC TGC GCC GCC GCC CGC CCG GCC CTC TTC CCC TCG GGC GTT CCT	144
Glu Pro Cys Ala Ala Ala Arg Pro Ala Leu Phe Pro Ser Gly Val Pro	
35 40 45	
GCG GCC CGG AGC AGC CCA GCG CAG CCC AGG CTT TGC CAA CGG CTC GAC	192
Ala Ala Arg Ser Ser Pro Ala Gln Pro Arg Leu Cys Gln Arg Leu Asp	
50 55 60	
GGG GCT TCT TGG GGA GTT TCT TA GGCCTGAAGG ACAAGAAGCA ACAACTCTGT	245
Gly Ala Ser Trp Gly Val Ser	
65 70	
TGATCAGAAC CTGTGGAAAA CCTCTGGCCT CTGTTTCAGAA TGAGTCCCAT GGGATTCCCC	305
GGCTGTGACA CTCTACCCTC CAGAACCTGA CGACTGGGCC ATGTGACCCA AGGAGGGATC	365
CTTACCAAGT GGGTTTTTAC CATCCTCTTG CTCTCTGTCT GAGAGATGTT TTCTAAACCC	425
CAGCCTTGAA CTTCACTCCT CCCTCAGTGG TAGTGTCCAG GTGCCGTGGA GCAGCAGGCT	485
GGCTTTGGTA GGGGCACCCA TCACCCGGCT TGCCTGTGCA GTCAGTGAGT GCTTAGGGCA	545
AAGAGAGCTC CCCTGGTTCC ATTCCTTCTG CCACCCAAAC CCTGATGAGA CCTTAGTGTT	605
CTCCAGGCTC TGTGGCCCAG GCTGAGAGCA GCAGGGTAGA AAAGACCAAG ATTGCGGGTT	665
TTATCTCTGG TTCCCTTATT ACTGCTCTCA AGCAGTGGCC TCTCTCACTT TAGCCATGGA	725
ATGGCTCCGA TCTACCTCAC AGCAGTGTCA GAAGGACTTC GCCAGGGTTT TGGGAGCTCC	785

SUBSTITUTE SHEET

-50-

AGGGTTCATA AGAAGGTGAA CCATTAGAAC AGATCCCTTC TTTTCCTTTT GCAATCAGAT 845
 AAATAAATAT CACTGAATGC AGTTC 870

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 71 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Asn Ser Ala Phe Asn Pro Leu Ile Tyr Cys Arg Ser Pro Asp Phe Arg
 1 5 10 15
 Ser Ala Phe Arg Arg Leu Leu Cys Arg Cys Gly Arg Arg Leu Pro Pro
 20 25 30
 Glu Pro Cys Ala Ala Ala Arg Pro Ala Leu Phe Pro Ser Gly Val Pro
 35 40 45
 Ala Ala Arg Ser Ser Pro Ala Gln Pro Arg Leu Cys Gln Arg Leu Asp
 50 55 60
 Gly Ala Ser Trp Gly Val Ser
 65 70

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2005 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cdna to mRNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 51..1250

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TAAGCCAGCG GGTCTGGGGG GAAAACTTCC CATCCCAGAC GCGACACGAG ATG GCT 56
 Met Ala
 1
 CCG TGG CCT CAC AAA AAC GGC TCT CTG GCT TTC TGG TCA GAC GCC CCC 104
 Pro Trp Pro His Lys Asn Gly Ser Leu Ala Phe Trp Ser Asp Ala Pro
 5 10 15
 ACC TTG GAC CCC AGT GCA GCC AAC ACC AGT GGG TTG CCA GGG GTG CCA 152
 Thr Leu Asp Pro Ser Ala Ala Asn Thr Ser Gly Leu Pro Gly Val Pro
 20 25 30
 TGG GCA GCG GCA TTG GCT GGA GCA TTG CTG GCG CTG GCC ACG GTG GGA 200
 Trp Ala Ala Ala Leu Ala Gly Ala Leu Leu Ala Leu Ala Thr Val Gly
 35 40 45 50

SUBSTITUTE SHEET

-51-

GGC AAC CTG CTG GTA ATC ACA GCT ATC GCC CGC ACG CCG AGA CTA CAG Gly Asn Leu Leu Val Ile Thr Ala Ile Ala Arg Thr Pro Arg Leu Gln 55 60 65	248
ACC ATA ACC AAC GTG TTC GTG ACT TCG CTG GCC ACA GCT GAC TTG GTA Thr Ile Thr Asn Val Phe Val Thr Ser Leu Ala Thr Ala Asp Leu Val 70 75 80	296
GTG GGA CTC CTC GTA ATG CCA CCA GGG GCC ACA TTG GCG CTG ACT GGC Val Gly Leu Leu Val Met Pro Pro Gly Ala Thr Leu Ala Leu Thr Gly 85 90 95	344
CAC TGG CCC TTG GGC GCA ACT GGC TGC GAG CTG TGG ACG TCA GTG GAC His Trp Pro Leu Gly Ala Thr Gly Cys Glu Leu Trp Thr Ser Val Asp 100 105 110	392
GTG CTC TGT GTA ACT GCC AGC ATC GAG ACC CTG TGC GCC CTG GCT GTA Val Leu Cys Val Thr Ala Ser Ile Glu Thr Leu Cys Ala Leu Ala Val 115 120 125 130	440
GAC CGC TAC CTA GCC GTC ACC AAC CCT CTG CGT TAC GGC ACG CTG GTT Asp Arg Tyr Leu Ala Val Thr Asn Pro Leu Arg Tyr Gly Thr Leu Val 135 140 145	488
ACC AAG CGC CGC GCC CGG GCG GCA GTA GTC CTG GTG TGG ATC GTG TCC Thr Lys Arg Arg Ala Arg Ala Ala Val Val Leu Val Trp Ile Val Ser 150 155 160	536
GCC ACC GTG TCC TTT GCG CCC ATC ATG AGC CAG TGG TGG CGT GTA GGG Ala Thr Val Ser Phe Ala Pro Ile Met Ser Gln Trp Trp Arg Val Gly 165 170 175	584
GCA GAC GCT GAG GCG CAA GAG TGT CAC TCC AAT CCG CGC TGC TGT TCC Ala Asp Ala Glu Ala Gln Glu Cys His Ser Asn Pro Arg Cys Cys Ser 180 185 190	632
TTT GCC TCC AAT ATG CCC TAC GCG CTG CTC TCC TCC TCC GTC TCC TTC Phe Ala Ser Asn Met Pro Tyr Ala Leu Leu Ser Ser Ser Val Ser Phe 195 200 205 210	680
TAC CTT CCC CTC CTT GTG ATG CTC TTC GTC TAT GCT CGA GTG TTC GTC Tyr Leu Pro Leu Leu Val Met Leu Phe Val Tyr Ala Arg Val Phe Val 215 220 225	728
GTA GCT AAG CGC CAG CGG CGT TTG CTG CGC CGG GAG CTG GGC CGT TTT Val Ala Lys Arg Gln Arg Arg Leu Leu Arg Arg Glu Leu Gly Arg Phe 230 235 240	776
CCG CCC GAG GAG TCT CCG CGG TCT CCG TCG CGC TCT CCA TCC CCT GCC Pro Pro Glu Glu Ser Pro Arg Ser Pro Ser Arg Ser Pro Ser Pro Ala 245 250 255	824
ACA GTC GGG ACA CCC ACG GCA TCG GAT GGA GTG CCC TCC TGC GGG CGG Thr Val Gly Thr Pro Thr Ala Ser Asp Gly Val Pro Ser Cys Gly Arg 260 265 270	872
CGG CCT GCG CGC CTC CTA CCG CTC GGG GAA CAC CGC GCC CTG CGC ACC Arg Pro Ala Arg Leu Leu Pro Leu Gly Glu His Arg Ala Leu Arg Thr 275 280 285 290	920
TTG GGT CTC ATT ATG GGC ATC TTC TCT CTG TGC TGG CTG CCC TTC TTT Leu Gly Leu Ile Met Gly Ile Phe Ser Leu Cys Trp Leu Pro Phe Phe 295 300 305	968

SEQUENCE SHEET

-52-

CTG GCC AAC GTG CTG CGC GCA CTC GTG GGG CCC TCC CTA GTT CCC AGC	1016
Leu Ala Asn Val Leu Arg Ala Leu Val Gly Pro Ser Leu Val Pro Ser	
310 315 320	
GGA GTT TTC ATC GCC CTG AAC TGG TTG GGC TAT GCC AAC TCT GCC TTC	1064
Gly Val Phe Ile Ala Leu Asn Trp Leu Gly Tyr Ala Asn Ser Ala Phe	
325 330 335	
AAC CCG CTC ATC TAC TGC CGC AGC CCG GAC TTT CGC GAC GCC TTC CGT	1112
Asn Pro Leu Ile Tyr Cys Arg Ser Pro Asp Phe Arg Asp Ala Phe Arg	
340 345 350	
CGT CTT CTG TGC AGC TAC GGT GGC CGT GGA CCG GAA GAG CCA CGC GTG	1160
Arg Leu Leu Cys Ser Tyr Gly Gly Arg Gly Pro Glu Glu Pro Arg Val	
355 360 365 370	
GTC ACC TTC CCA GCT AGC CCT GTT GCG TCC AGG CAG AAC TCA CCG CTC	1208
Val Thr Phe Pro Ala Ser Pro Val Ala Ser Arg Gln Asn Ser Pro Leu	
375 380 385	
AAC AGG TTT GAT GGC TAT GAA GGT GAG CGT CCA TTT CCC ACA	1250
Asn Arg Phe Asp Gly Tyr Glu Gly Glu Arg Pro Phe Pro Thr	
390 395 400	
TGAAGGACCA TGGAGATCTA GCAAGGAGCC TGACTTCTGG AGAAATTTTT TTTTAAGACA	1310
GAAAGACAAG CAACGTCCAT GGATGCAAAC CTTTATCAG CCCTTGATTC TGCTCAGAGT	1370
GAGTTCCAG GAACCGCAAC TCTCCAGACC ATGCATAGAC CACAGAATGT AAAGGGGAAA	1430
TCTTACCAA TGGGTTTACC ATCTTCTCTC TCTTCGTGAG AGTGTCTATA GGCCACCTTG	1490
AACTTCGCTA CTACCTCAGC CGCCGGATAT CAGCCACCCT GCGTTGACTG CCTGGGAGGA	1550
GCTGCGTTCC CACCACCACC CTGCTTATTA TGTTTGTGCT GGATGCTTAG GGCTAAGAAA	1610
GCACCCTTAC CTACCTCCCT TCCTACGCTT TCCTGACCCC ATGAATGACT TTTGTCTCCA	1670
CAAATCACTC TGTCTCCAGG TTCTGTGTTT CCAGTCTCTG TGTCTCTGGT TAGTTGGAAA	1730
GCAGGAAACC CGCGGGGGGA GCGGGGGGAG GGGGGGAACG ACCAAGTTTG AGGTTTTGTG	1790
CCTGGCTCCT CACTACAGCT CTCTAACAT CATCTTGGAC CATCTCTCAC AATAGGCACA	1850
AAACAGCTCT AATCTACCTC ACTCTTAGGA CTTCAAGGTT TGGGAGAAAT TCCAGGGTTC	1910
CTGGGAAGAA GTCAAACCAT TGGAAATGGGT CCCTTTTGGC GTTAAATCA AATTAATAAA	1970
TATTATTGAA TGTGAAAAAA AAAAAAAAT CTAGA	2005

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 400 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Ala Pro Trp Pro His Lys Asn Gly Ser Leu Ala Phe Trp Ser Asp
 1 5 10 15

SUBSTITUTE SHEET

-53-

Ala Pro Thr Leu Asp Pro Ser Ala Ala Asn Thr Ser Gly Leu Pro Gly
 20 25 30
 Val Pro Trp Ala Ala Ala Leu Ala Gly Ala Leu Leu Ala Leu Ala Thr
 35 40 45
 Val Gly Gly Asn Leu Leu Val Ile Thr Ala Ile Ala Arg Thr Pro Arg
 50 55 60
 Leu Gln Thr Ile Thr Asn Val Phe Val Thr Ser Leu Ala Thr Ala Asp
 65 70 75 80
 Leu Val Val Gly Leu Leu Val Met Pro Pro Gly Ala Thr Leu Ala Leu
 85 90 95
 Thr Gly His Trp Pro Leu Gly Ala Thr Gly Cys Glu Leu Trp Thr Ser
 100 105 110
 Val Asp Val Leu Cys Val Thr Ala Ser Ile Glu Thr Leu Cys Ala Leu
 115 120 125
 Ala Val Asp Arg Tyr Leu Ala Val Thr Asn Pro Leu Arg Tyr Gly Thr
 130 135 140
 Leu Val Thr Lys Arg Arg Ala Arg Ala Ala Val Val Leu Val Trp Ile
 145 150 155 160
 Val Ser Ala Thr Val Ser Phe Ala Pro Ile Met Ser Gln Trp Trp Arg
 165 170 175
 Val Gly Ala Asp Ala Glu Ala Gln Glu Cys His Ser Asn Pro Arg Cys
 180 185 190
 Cys Ser Phe Ala Ser Asn Met Pro Tyr Ala Leu Leu Ser Ser Ser Val
 195 200 205
 Ser Phe Tyr Leu Pro Leu Leu Val Met Leu Phe Val Tyr Ala Arg Val
 210 215 220
 Phe Val Val Ala Lys Arg Gln Arg Arg Leu Leu Arg Arg Glu Leu Gly
 225 230 235 240
 Arg Phe Pro Pro Glu Glu Ser Pro Arg Ser Pro Ser Arg Ser Pro Ser
 245 250 255
 Pro Ala Thr Val Gly Thr Pro Thr Ala Ser Asp Gly Val Pro Ser Cys
 260 265 270
 Gly Arg Arg Pro Ala Arg Leu Leu Pro Leu Gly Glu His Arg Ala Leu
 275 280 285
 Arg Thr Leu Gly Leu Ile Met Gly Ile Phe Ser Leu Cys Trp Leu Pro
 290 295 300
 Phe Phe Leu Ala Asn Val Leu Arg Ala Leu Val Gly Pro Ser Leu Val
 305 310 315 320
 Pro Ser Gly Val Phe Ile Ala Leu Asn Trp Leu Gly Tyr Ala Asn Ser
 325 330 335
 Ala Phe Asn Pro Leu Ile Tyr Cys Arg Ser Pro Asp Phe Arg Asp Ala
 340 345 350

SUBSTITUTE SHEET

-54-

Phe Arg Arg Leu Leu Cys Ser Tyr Gly Gly Arg Gly Pro Glu Glu Pro
 355 360 365

Arg Val Val Thr Phe Pro Ala Ser Pro Val Ala Ser Arg Gln Asn Ser
 370 375 380

Pro Leu Asn Arg Phe Asp Gly Tyr Glu Gly Glu Arg Pro Phe Pro Thr
 385 390 395 400

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 687 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

- (ix) FEATURE:
 (A) NAME/KEY: intron
 (B) LOCATION: 9..402

- (ix) FEATURE:
 (A) NAME/KEY: exon
 (B) LOCATION: 403..470

- (ix) FEATURE:
 (A) NAME/KEY: intron
 (B) LOCATION: 471..674

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CTCAACAGGT AGCGGACGCA GGCAGAGGAC TGGAGTCTGG GTGGGGACGC CTCTGTCTCT	60
ATTTTGTAGT TTGAGGGTTG GGGGAGGAGA AGGTGTAGAC AGGGCTTTTG TCTCGAGAGG	120
ACAGAAAAGG AGTAAGAACA GAATCGGGAT CTAGGGCCCT TCCTTTTATT GGATCCAATC	180
CCTGGGTCTG AGGCAAAGGA GGAAAGGGAA ATTTGTTTAC CTTGGGACCA GGTGAGCCCC	240
ACAGGTTTCT GCCAGCAGGT TTCTGACCTC TCTGGTTGCC TCTAGTTTGG ATCTTTTATG	300
TTCTATTCTC CAGGCGCCCA GGTATCACTA ACTTGTCTGG GACATCCATA GACAGCAATG	360
GACATGTCAA GTCTCTGCC TCAGTTCCGC TTTCTTTCAA AGGTTTGATG GCTATGAAGG	420
TGAGCGTCCA TTTCCACAT GAAGGACCAT GGAGATCTAG CAAGGAGCCT GTGAGTTGAA	480
TTTGAGCTGC TTTTCTCCCT CAGGGACTGG ATTCGAGGTG TAGGGTGGGA TGAGGGAGGG	540
TGCAGGATGA TCCCTATATC TTTGAAAAGT AAATATGCTA TTCAGGGTTC CTGAGTCACT	600
CCCCTCTTAC CTCCAGTGCT TGATCCGCAC CTCCTTGACT GGTTACCCCA AGAAATATTG	660
TTTCCGTTTT GCAGGACTTC TGGAGAA	687

SUBSTITUTE SHEET

-55-

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 176 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

- (ix) FEATURE:
 (A) NAME/KEY: exon
 (B) LOCATION: 1..60

- (ix) FEATURE:
 (A) NAME/KEY: exon
 (B) LOCATION: 61..176

- (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 2..97

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

A CGC GCA GTC ACC TTC CCA GCC AGC CCT GTT GAA GCC AGG CAG AGT	46
Arg Ala Val Thr Phe Pro Ala Ser Pro Val Glu Ala Arg Gln Ser	
1 5 10 15	
CCA CCG CTC AAC AGG TTT GAT GGC TAT GAA GGT GCG CGT CCG TTT CCC	94
Pro Pro Leu Asn Arg Phe Asp Gly Tyr Glu Gly Ala Arg Pro Phe Pro	
20 25 30	
ACG TGAAGGGCCG TGAAGATCCA GCAAGGAAGC TGACTTCTGG GGATTTTTTT	147
Thr	
TTTCCTCCAG AAAGACAAGC AACGTCCAT	176

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Arg Ala Val Thr Phe Pro Ala Ser Pro Val Glu Ala Arg Gln Ser Pro	
1 5 10 15	
Pro Leu Asn Arg Phe Asp Gly Tyr Glu Gly Ala Arg Pro Phe Pro Thr	
20 25 30	

SUBSTITUTE SHEET

-56-

CLAIMS

- 5 1. A DNA sequence which encodes a
mammalian β_3 -adrenergic receptor in substantially
pure form, functional equivalents thereof or a
nucleic acid sequence which hybridizes thereto.
- 10 2. The DNA sequence of claim 1 which
encodes human β_3 -adrenergic receptor.
3. The DNA sequence of claim 2 which
comprises genomic DNA in a combination with cDNA.
- 15 4. The gene of claim 2 which comprises
the coding sequence depicted in Fig. 1 hereof.
5. A nucleic acid sequence of claim 1
20 which is cDNA.
6. A nucleic acid sequence of claim 5
which is p184 depicted in Fig. 10B hereof.
- 25 7. A nucleic acid sequence of claim 5
which is p192 described herein.

SUBSTITUTE SHEET

-57-

8. A nucleic acid sequence of claim 1
which is mRNA or cRNA.

5 9. The gene of claim 1 which encodes
rodent β_3 -adrenergic receptor.

10. The gene of claim 9 which encodes
rat β_3 -adrenergic receptor.

10

11. An oligonucleotide which
hybridizes to the DNA sequence of claim 1.

12. An oligonucleotide which
15 hybridizes to a nucleic acid of claim 1.

13. An oligonucleotide of claim 11
which carries a detectable label.

20 14. A vector which comprises the DNA
of claim 1.

15. A host cell transformed with the
vector of claim 14.

25

SUBSTITUTE SHEET

-58-

16. A vector which comprises the DNA
of claim 2.

17. A host cell which is transformed
5 with the vector of claim 16.

18. A host cell of claim 17 which does
not express other β adrenergic receptors.

10 19. A vector which comprises the DNA
of claim 4.

20. A vector of claim 19 which is a
shuttle vector.

15

21. A host cell which is transformed
with the vector of claim 20.

22. A host cell of claim 21 which does
20 not express other β -adrenergic receptors

23. A method of preparing a β_3 -
adrenergic receptor which comprises culturing a
host cell of claim 18.

25

SUBSTITUTE SHEET

-59-

24. A method of preparing a β_3 -adrenergic receptor which comprises culturing a host cell of claim 22.

5 25. A method for monitoring the presence of human β_3 receptor gene which comprises extracting the mRNA from human adipose tissue and bringing said mRNA into contact with a nucleic acid sequence which hybridizes to the DNA
10 sequence of claim 2.

26. The method of claim 25 which utilizes the nucleic acid sequence of claim 7.

15 27. The method of claim 26 wherein the mRNA hybridizes to the nucleic acid sequence of claim 6.

28. The method of claim 26 wherein the
20 mRNA hybridizes to the nucleic acid sequence of claim 9.

29. A method for identifying a compound which affect the activity of the β_3 -
25 adrenergic receptor which comprises bringing said compound in contact with a host cell transformed

SUBSTITUTE SHEET

-60-

with a vector of claim 14 and detecting any change in the level of activity of the β_3 -adrenergic receptor.

5 30. The method of claim 29 wherein the host cell is transformed with the vector of claim 16.

10 31. The method of claim 30 wherein the host cell does not express other β -adrenergic receptors.

15 32. The method of claim 29 wherein the host cell is transformed with the vector of claim 19.

20 33. The method claim 32 wherein the host cell does not express other β -adrenergic receptors.

25 34. A β_3 -adrenergic receptor protein in substantially pure form having the amino acid sequence depicted in Fig. 1 and functional equivalents thereof.

SUBSTITUTE SHEET

-61-

35. A β_3 -adrenergic receptor protein
produced by the expression of the DNA of claim 1.

36. The receptor protein of claim 35
5 which is human β_3 -adrenergic receptor protein.

37. The receptor protein of claim 36
which is rodent β_3 -adrenergic receptor protein.

10 38. A DNA sequence which encodes the
first exon of β_3 -adrenergic receptor wherein the
nucleotide at position 1206 is changed from
guanine (G) to a nucleotide selected from thymine
(T), adenine (A) or cytosine (C).

15 39. The DNA sequence of claim 38 which
encodes the first exon of human β_3 -adrenergic
receptor.

20 40. An antibody which is specific to
the protein of claim 34.

41. The antibody of claim 40 which is
a monoclonal antibody.

25

SUBSTITUTE SHEET

-62-

42. An antibody which is specific to the protein of claim 36.

43. The antibody of claim 42 which is
5 a monoclonal antibody.

44. A DNA construct which comprises the fat-specific promoter and enhancer elements of the rodent β_3 -adrenergic receptor.

10

45. The construct of claim 44 wherein the DNA sequence is selected from the introns and the 5'-flanking region.

15

46. The construct of claim 45 which is a vector.

47. The construct of claim 46 which contains a reporter gene whose transcription is
20 modulated by the fat-specific promoter and enhancer elements of the rat β_3 receptor gene.

48. A host cell transfected with the construct of claim 47.

25

SUBSTITUTE SHEET

-63-

49. The host cell of claim 48 wherein the fat-specific DNA sequences are from the 5'-flanking region of the rat β_3 receptor gene.

5 50. The host cell of claim 48 wherein the fat-specific DNA sequences are from the introns of the rat β_3 receptor gene.

SUBSTITUTE SHEET

1/12

			10				20				30				40			
ATG	GCT	CCG	TGG	CCT	CAC	GAG	AAC	AGC	TCT	CTT	GCC	CCA	TGG	CCG				
MET	Ala	Pro	Trp	Pro	His	Glu	Asn	Ser	Ser	Leu	Ala	Pro	Trp	Pro				
			50				60				70				80			90
GAC	CTC	CCC	ACC	CTG	GCG	CCC	AAT	ACC	GCC	AAC	ACC	AGT	GGG	CTG				
Asp	Leu	Pro	Thr	Leu	Ala	Pro	Asn	Thr	Ala	Asn	Thr	Ser	Gly	Leu				
			100				110				120				130			
CCA	GGG	GTT	CCG	TGG	GAG	GCG	GCC	CTA	GCC	GGG	GCC	CTG	CTG	GCG				
Pro	Gly	Val	Pro	Trp	Glu	Ala	Ala	Leu	Ala	Gly	Ala	Leu	Leu	Ala				
			140				150				160				170			180
CTG	GCG	GTG	CTG	GCC	ACC	GTG	GGA	GGC	AAC	CTG	CTG	GTC	ATC	GTG				
Leu	Ala	Val	Leu	Ala	Thr	Val	Gly	Gly	Asn	Leu	Leu	Val	Ile	Val				
			190				200				210				220			
GCC	ATC	GCC	TGG	ACT	CCG	AGA	CTC	CAG	ACC	ATG	ACC	AAC	GTG	TTC				
Ala	Ile	Ala	Trp	Thr	Pro	Arg	Leu	Gln	Thr	MET	Thr	Asn	Val	Phe				
			230				240				250				260			270
GTG	ACT	TCG	CTG	GCC	GCA	GCC	GAC	CTG	GTG	ATG	GGA	CTC	CTG	GTG				
Val	Thr	Ser	Leu	Ala	Ala	Ala	Asp	Leu	Val	MET	Gly	Leu	Leu	Val				
			280				290				300				310			
GTG	CCG	CCG	GCG	GCC	ACC	TTG	GCG	CTG	ACT	GCG	CAC	TGG	CCG	TTG				
Val	Pro	Pro	Ala	Ala	Thr	Leu	Ala	Leu	Thr	Gly	His	Trp	Pro	Leu				
			320				330				340				350			360
GGC	GCC	ACT	GCG	TGC	GAG	CTG	TGG	ACC	TCG	GTG	GAC	GTG	CTG	TGT				
Gly	Ala	Thr	Gly	Cys	Glu	Leu	Trp	Thr	Ser	Val	Asp	Val	Leu	Cys				

Fig-1A

SUBSTITUTE SHEET

2/12

			370				380				390				400			
GTG	ACC	GCC	AGC	ATC	GAA	ACC	CTG	TGC	GCC	CTG	GCC	GTG	GAC	CGC				
Val	Thr	Ala	Ser	Ile	Glu	Thr	Leu	Cys	Ala	Leu	Ala	Val	Asp	Arg				
			410				420				430				440			450
TAC	CTG	GCT	GTG	ACC	AAC	CCG	CTG	CGT	TAC	GGC	GCA	CTG	GTC	ACC				
Tyr	Leu	Ala	Val	Thr	Asn	Pro	Leu	Arg	Tyr	Gly	Ala	Leu	Val	Thr				
			460				470				480				490			
AAG	CGC	TGC	GCC	CGG	ACA	GCT	GTG	GTC	CTG	GTG	TGG	GTC	GTG	TCG				
Lys	Arg	Cys	Ala	Arg	Thr	Ala	Val	Val	Leu	Val	Trp	Val	Val	Ser				
			500				510				520				530			540
GCC	GCG	GTG	TCG	TTT	GCG	CCC	ATC	ATG	AGC	CAG	TGG	TGG	CGC	GTA				
Ala	Ala	Val	Ser	Phe	Ala	Pro	Ile	MET	Ser	Gln	Trp	Trp	Arg	Val				
			550				560				570				580			
GGG	GCC	GAC	GCC	GAG	GCG	CAG	CGC	TGC	CAC	TCC	AAC	CCG	CGC	TGC				
Gly	Ala	Asp	Ala	Glu	Ala	Gln	Arg	Cys	His	Ser	Asn	Pro	Arg	Cys				
			590				600				610				620			630
TGT	GCC	TTC	GCC	TCC	AAC	ATG	CCC	TAC	GTG	CTG	CTG	TCC	TCC	TCC				
Cys	Ala	Phe	Ala	Ser	Asn	MET	Pro	Tyr	Val	Leu	Leu	Ser	Ser	Ser				
			640				650				660				670			
GTC	TCC	TTC	TAC	CTT	CCT	CTT	CTC	GTG	ATG	CTC	TTC	GTC	TAC	GCG				
Val	Ser	Phe	Tyr	Leu	Pro	Leu	Leu	Val	MET	Leu	Phe	Val	Tyr	Ala				
			680				690				700				710			720
CGG	GTT	TTC	GTG	GTG	GCT	ACG	CGC	CAG	CTG	CGC	TTG	CTG	CGC	GGG				
Arg	Val	Phe	Val	Val	Ala	Thr	Arg	Gln	Leu	Arg	Leu	Leu	Arg	Gly				
			730				740				750				760			
GAG	CTG	GCG	CGC	TTT	CCG	CCC	GAG	GAG	TCT	CCG	CCG	GCG	CCG	TCG				
Glu	Leu	Gly	Arg	Phe	Pro	Pro	Glu	Glu	Ser	Pro	Pro	Ala	Pro	Ser				

Fig-1B

SUBSTITUTE SHEET

3/12

770	780	790	800	810
CGC TCT CTG GCC CCG GCC CCG GTG GGG ACG TGC GCT CCG CCC GAA				
Arg Ser Leu Ala Pro Ala Pro Val Gly Thr Cys Ala Pro Pro Glu				
820	830	840	850	
GGG GTG CCC GCC TGC GGC CGG CGG CCC GCG CGC CTC CTG CCT CTC				
Gly Val Pro Ala Cys Gly Arg Arg Pro Ala Arg Leu Leu Pro Leu				
860	870	880	890	900
CGG GAA CAC CGG GCC CTG TGC ACC TTG GGT CTC ATC ATG GGC ACC				
Arg Glu His Arg Ala Leu Cys Thr Leu Gly Leu Ile MET Gly Thr				
910	920	930	940	
TTC ACT CTC TGC TGG TTG CCC TTC TTT CTG GCC AAC GTG CTG CGC				
Phe Thr Leu Cys Trp Leu Pro Phe Phe Leu Ala Asn Val Leu Arg				
950	960	970	980	990
GCC CTG GGG GGC CCC TCT CTA GTC CCG GGC CCG GCT TTC CTT GCC				
Ala Leu Gly Gly Pro Ser Leu Val Pro Gly Pro Ala Phe Leu Ala				
1000	1010	1020	1030	
CTG AAC TGG CTA GGT TAT GCC AAT TCT GCC TTC AAC CCG CTC ATC				
Leu Asn Trp Leu Gly Tyr Ala Asn Ser Ala Phe Asn Pro Leu Ile				
1040	1050	1060	1070	1080
TAC TGC CGC AGC CCG GAC TTT CGC AGC GCC TTC CGC CGT CTT CTG				
Tyr Cys Arg Ser Pro Asp Phe Arg Ser Ala Phe Arg Arg Leu Leu				
1090	1100	1110	1120	
TGC CGC TGC GGC CGT CGC CTG CCT CCG GAG CCC TGC GCC GCC GCC				
Cys Arg Cys Gly Arg Arg Leu Pro Pro Glu Pro Cys Ala Ala Ala				
1130	1140	1150	1160	1170
CGC CCG GCC CTC TTC CCC TCG GGC GTT CCT GCG GCC CGG AGC AGC				
Arg Pro Ala Leu Phe Pro Ser Gly Val Pro Ala Ala Arg Ser Ser				

Fig-1C

SUBSTITUTE SHEET

4/12

			1180				1190				1200				1210		
CCA	GCG	CAG	CCC	AGG	CTT	TGC	CAA	CGG	CTC	GAC	GGG	GCT	TCT	TGG			
Pro	Ala	Gln	Pro	Arg	Leu	Cys	Gln	Arg	Leu	Asp	Gly	Ala	Ser	Trp			
			1220														
			GGA	GTT	TCT	TAG											
			Gly	Val	Ser	---											

Fig-1D

SUBSTITUTE SHEET

5/12




Fig-2A




Fig-2B




Fig-3A




Fig-3B




Fig-4

7/12

	GENOMIC	CDNA
Rat	CCG CTC AAC <u>AGG TAG</u> Pro Leu Asn Arg	CCG CTC AAC AGG TTT GAT GGC TAT GAA Pro Leu Asn Arg Phe Asp Gly Tyr Glu GGT GAG CGT CCA TTT CCC ACA TGA Gly Glu Arg Pro Phe Pro Thr <u>STOP</u>
Mouse	CCG CTC AAC <u>AGG TAG</u> Pro Leu Asn Arg	
Human	CGG CTC GAC <u>GGG TAG</u> Arg Leu Asp Gly	

Fig-5

...GTT GAA GCC AGG CAG AGT CCA CCG CTC AAC AGG TTT GAT GGC TAT
Val Glu Ala Arg Gln Ser Pro Pro Leu Asp Arg Phe Asp Gly Tyr

GAA GGT GCG CGT CCG TTT CCC ACG TGA AGGGCCGTGAAGATCCAGCAAG
Glu Gly Ala Arg Pro Phe Pro Thr ---

GAAGCTGACTTCTGGGGATTTTTTTTTTTCCTCCAGAAAGACAAGCAACGTCCAT...

Fig-6

SUBSTITUTE SHEET




Fig-7




Fig-8

9/12




Fig-9A




Fig-9B

SUBSTITUTE SHEET

10/12

(162 bp to begining of clone)... GCG CAG CCC AGG CTT TGC CAA
Ala Gln Pro Arg Leu Cys Gln

CGG CTC GAC GGG GCT TCT TGG GGA GTT TCT TAG GCCTGAAGGACAAGAA
Arg Leu Asp Gly Ala Ser Trp Gly Val Ser ---

GCAACAACCTCTGTTGATCAGAACCTGTGGAAA...(680 bp to poly A)

Fig-10A

AATTCTGCCT	TCAACCCGCT	CATCTACTGC	CGCAGCCCGG	ACTTTCGCAG
CGCCTTCCGC	CGTCTTCTGT	GCCGCTGCGG	CCGTCGCCTG	CCTCCGGAGC
CCTGCGCCGC	CGCCCGCCCG	GCCCTCTTCC	CCTCGGGCGT	TCCTGCGGCC
CGGAGCAGCC	CAGCGCAGCC	CAGGCTTTGC	CAACGGCTCG	ACGGGGCTTC
TTGGGGAGTT	TCTTAGGCCT	GAAGGACAAG	AAGCAACAAC	TCTGTTGATC
AGAACCTGTG	GAAAACCTCT	GGCCTCTGTT	CAGAATGAGT	CCCATGGGAT
TCCCCGGCTG	TGACACTCTA	CCCTCCAGAA	CCTGACGACT	GGGCCATGTG
ACCCAAGGAG	GGATCCTTAC	CAAGTGGGTT	TTCACCATCC	TCTTGCTCTC
TGTCTGAGAG	ATGTTTTCTA	AACCCAGCC	TTGAACTTCA	CTCCTCCCTC
AGTGGTAGTG	TCCAGGTGCC	GTGGAGCAGC	AGGCTGGCTT	TGGTAGGGGC
ACCCATCACC	CGGCTTGCCCT	GTGCAGTCAG	TGAGTGCTTA	GGGCAAAGAG
AGCTCCCCCTG	GTTCCATTCC	TTCTGCCACC	CAAACCCTGA	TGAGACCTTA
GTGTTCTCCA	GGCTCTGTGG	CCCAGGCTGA	GAGCAGCAGG	GTAGAAAAGA
CCAAGATTTG	GGGTTTTATC	TCTGGTTCCC	TTATTACTGC	TCTCAAGCAG
TGGCCTCTCT	CACTTTAGCC	ATGGAATGGC	TCCGATCTAC	CTCACAGCAG
TGTCAGAAGG	ACTTCGCCAG	GGTTTTGGGA	GCTCCAGGGT	TCATAAGAAG
GTGAACCATT	AGAACAGATC	CCTTCTTTTC	CTTTTGCAAT	CAGATAAATA
AATATCACTG	AATGCAGTTC			

Fig-10B

SUBSTITUTE SHEET




Fig-11




Fig-12




Fig-13

12/12




Fig-14

SUBSTITUTE SHEET

INTERNATIONAL SEARCH REPORT

national application No.
PCT/US93/06733

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C12N 1/21, 15/00; C07K 13/00, 15/28
US CL : 536/23.1; 530/350, 388.1; 435/69.1, 240.1, 320.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.1; 530/350, 388.1; 435/69.1, 240.1, 320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	WO, A, 90/08775 (Emorine, et al) 09 August 1990, see abstract.	1-3, 5, 8-18, 23, 25, 27-31, 34, 35, 36, 40-43 37, 44-50; 4, 6, 7, 19-22, 24, 26, 32, 33, 38, 39

<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.	
<p>* Special categories of cited documents</p> <p>"A" document defining the general state of the art which is not considered to be part of particular relevance</p> <p>"E" earlier document published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"A" document member of the same patent family</p>
Date of the actual completion of the international search 23 SEPTEMBER 1993	Date of mailing of the international search report OCT 06 1993
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. NOT APPLICABLE	Authorized officer ROBERT HILL <i>Robert Hill</i> Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/06733

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	Science, Volume 245, issued 08 September 1989, L. J. Emorine et al, "Molecular Characterization of the Human B3-Adrenergic Receptor", pages 1118-1121, especially figure 1.	1-3, 5, 8-18, 23, 25, 27-31, 34, <u>35, 36, 40-43</u> , 37, 44-50, 4, 6, 7, 19-22, 24, 26, 32, 33, 38, 39
X Y	Molecular Pharmacology, Volume 40, issued 1991, J. G. Granneman et al, "Molecular Cloning and Expression of the Rat B3-Adrenergic Receptor", pages 895-899, especially figures 1 and 2.	1-3, 5, 8-18, 23, 25, 27-31, 34-36, <u>37, 40-43, 44-50</u> , 4, 6, 7, 19-22, 24, 26, 32, 33, 38, 39
X Y	The EMBO Journal, Volume 10, No. 12, issued 1991, C. Nahmias et al, "Molecular Characterization of the Mouse B3-Adrenergic Receptor: Relationship with the Atypical Receptor of Adipocytes", pages 3721-3727, especially figure 1.	1-3, 5, 8-18, 23, 25, 27-31, 34-36, <u>37, 40-43, 44-50</u> , 4, 6, 7, 19-22, 24, 26, 32, 33, 38, 39

INTERNATIONAL SEARCH REPORT

ational application No.

PCT/US93/06733

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, Medline, Dialog

search terms: B3-adrenergic receptor, cloning, cDNA, adipocyte receptor

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE LEFT BLANK